

Mid-air collisions enhance saltation

M. V. Carneiro,^{1,*} N. A. M. Araújo,^{2,3,†} T. Pähz,^{2,3,‡} and H. J. Herrmann^{1,4,§}

¹*Institut für Baustoffe, ETH-Hönggerberg, Schafmattstrasse 6, 8093 Zürich, Switzerland*

²*Department of Ocean Science and Engineering, Zhejiang University, 310058 Hangzhou, China*

³*State Key Laboratory of Satellite Ocean Environment Dynamics,
Second Institute of Oceanography, Hangzhou, China*

⁴*Departamento de Física, Universidade Federal do Ceará, 60451-970 Fortaleza, Ceará, Brazil*

Here we address the old question in Aeolian particle transport about the importance of mid-air collisions. We find that surprisingly these collisions do enhance the overall flux substantially. The effect depends strongly on restitution coefficient and wind speed. We can explain this observation as a consequence of a soft bed of grains which floats above the surface and reflects the highest flying particles. We make the unexpected observation that the flux is maximized for an intermediate restitution coefficient of about 0.65, which is comparable to the values experimentally measured for collisions between sand grains.

PACS numbers: 45.70.Mg, 47.27.-i, 47.55.Kf, 83.80.Hj

Would a sandstorm be stronger if the sand grains in the air would not collide against each other? This question has puzzled practitioners and theoreticians alike in the past. Models for Aeolian sand flux in Refs. [1–6] become certainly much simpler if such mid-air collisions are neglected, but does this approximation underestimate or overestimate the value of the calculated saturated flux? As opposed to experiments, the direct computer simulation of saltation, the main Aeolian transport process, offers the possibility of switching on or off the collisions between particles or of modifying at will the collision parameters, like for instance the coefficient of restitution, allowing in this way to determine for the first time with precision the role of mid-air collisions during the saltation process.

We discover that mid-air collisions are in fact the key ingredient to understand the relation between different concepts that have been put forward in the past like the splash [7, 8], the soft bed [9, 10], and the distinction between saltans and reptons [11, 12]. In saltation, particles are ejected from the granular bed in a splash, produced by the impact of fast particles called saltans (yellow trajectory in Fig. 1). These saltans must have the necessary kinetic energy to assure that, despite the substantial dissipation in the ground, some ejected particles can again fly high enough. After all, only high-flyers can acquire sufficient acceleration to become again saltans, because the wind velocity at the ground is zero, increasing logarithmically with height. Our detailed study reveals the following picture: Due to the irregularities of the surface a splash produces in fact three types of moving particles (see Fig. 1): Many (green) creepers that do not leave the bed, many (red) leapers making small jumps remaining in regions of small wind velocities and that, therefore, cannot produce a new splash, and very few saltans (yellow) which fly higher up. Only the saltans assure that the saltation process is sustained. Both creepers and leapers (reptons) contribute considerably to the global sand flux




FIG. 1. (color online). Typical splash obtained numerically with our model in 3D. The impinging particle (yellow) bounces after ejecting other particles from the bed (red and green). While the green particles essentially only move on the ground, the red ones are lifted and dragged by the wind.

but play a very different role in what follows.

Can one really make such a sharp distinction between leapers and saltans? After all, the collision process is random and should yield a continuous distribution of ejection velocities. The experimental observations say yes: Already Mitha et al. [13] observed that the splash distribution is bimodal as later reproduced numerically by Anderson and Haff [14], i.e., that there is a broad spectrum of slow particles and a small peak of fast ones, and the analysis of these experimental observations led Andreotti [11] to coin the terms saltans and reptons. What makes these two types of particles different are the mid-air collisions. In Fig. 2, we see the trajectory of a typical saltan simulated with the method described below in three dimension: It makes several jumps without touching the ground, rebounding upwards each time due to a collision with a (red) leaper. The probability for such a collision is reasonably high because the leapers form a rather dense layer which is precisely the soft bed described earlier in Ref. [10]. Consequently, the saltan stays longer in areas




FIG. 2. (color online). Simulated trajectory of a salton in three dimensional saltation. The (yellow, upper trajectory) salton is kept in the air by colliding against (red, lower trajectories) particles from the soft bed with $e = 0.7$ and for Shields number $\theta = 0.9$. For clarity only the relevant particle trajectories are shown and the ground is schematically represented by a flat plane. (details in the text).

of strong wind and less time close to the ground, where the wind drag acceleration is weaker. This explains why the saltons acquire so much energy and can sustain the saltation process. Summarizing, mid-air collisions in the soft bed are the crucial mechanism that differentiates the saltons, essentially doubling the saturated sand flux as we will show in the following.

The Discrete Elements Method (DEM) based on particle-particle and particle-wind interactions allowing to study quantitatively, the Aeolian transport of granular systems [15]. It has been used to calculate the onset of saltation and confirm the existence of a jump in the total flux in two dimensions [16]. Here, we apply a three dimensional DEM scheme to investigate the role of mid-air collisions in saltation.

Every sand grain is represented by a hard sphere of average diameter D_{mean} . The collisions between particles are described in detail in the Supplemental Material [17]. We distinguish collisions between particles in the bed with restitution coefficient $e_{bed} = 0.7$ and between flying particles (mid-air collisions) with restitution coefficient e . Gravity acts in vertical direction (y -direction) and a logarithmic wind velocity profile $u(y)$ is imposed in the horizontal direction (x -direction),

$$u(y) = \frac{u_*}{\kappa} \ln \frac{y - h_0}{y_0}, \quad (1)$$

where $y_0 = D_{mean}/30$ is the roughness of the bed, h_0 the bed height, $\kappa = 0.4$ the von Kármán constant, and u_* the wind shear velocity. The Shields number defined as

$$\theta = \frac{u_*^2}{(s-1)gD_{mean}} \quad (2)$$

is the pertinent dimensionless parameter that controls the wind velocity, where $s = \rho_s/\rho_w$ is the ratio between the grain and fluid density, and g the norm of the gravitational acceleration. The feedback procedure that extracts the momentum from the wind due to the acceleration of




FIG. 3. (color online). Concentration and saturated flux. (a) Concentration profile of particles as a function of the height y for $\theta = 0.9$ in the absence (red, lower curve) and the presence (yellow, upper curve) of mid-air collisions ($e = 0.7$). (b) The relation between the saturated flux and the restitution coefficient, for three different Shields numbers exhibits a peak around $e = 0.65$ for higher shear velocities. The arrows show the flux without mid-air collisions.

the grains is also explained in the Supplemental Material [17].

We consider particles of average diameter $D_{mean} = 2 \times 10^{-4}$ m, size dispersion $\sigma_D = 0.15D_{mean}$, and density $\rho_s = 2650$ kg/m³, in a three-dimensional wind channel of dimension $(700 \times 75 \times 7.5)D_{mean}^3$ with a reflective upper boundary, placed sufficiently high to avoid any particle colliding against it, and periodic boundaries in the other directions. For the fluid density we considered $\rho_s = 1.174$ kg/m³. In fact, no collision with the upper boundary ever occurred in our simulations. The lower boundary at $y = 0$, representing the deep ground, is strongly dissipative with a fixed restitution coefficient of $e_w = 0.5$. We consider a bed of 12 particle layers to suppress the reflection of shock waves on the lower boundary due to the finite depth [18–20]. In the beginning of the simulation, a few particles are dropped at random positions to trigger saltation.

The dimensionless flux in the direction of the wind is defined as,

$$q = \frac{1}{DA} \sum_i^N m_i v_i^x, \quad (3)$$

where $A = (75 \times 7.5)D_{mean}^2$ is the area of the bottom of the channel, v_i^x and m_i are, respectively, the velocity along the horizontal direction and the mass of the particle i , and $D = \rho_s \sqrt{(s-1)gD_{mean}^3}$ is a normalization constant. The saturated flux, which is the average flux in the stationary state, does not change with the number of particles in the system. The granular temperature




FIG. 4. (color online) The temperature (a) and the flux (b) profiles for $e = 0.7$, 1.0 , and without mid-air collisions, for $\theta = 0.9$. At every height in (a), the granular temperature for $e = 1.0$ is larger than $e = 0.7$. The flux profiles in (b) confirm the higher flux for $e = 0.7$. The flux profile is defined at each height as the product of the concentration and the average velocity in each slice. (c) The dependence of the maximum temperature on the Shields number indicates a linear and a sublinear growth for the conservative and dissipative case, respectively.

defined as

$$T(y) = \frac{1}{N(y)} \sum_i^{N(y)} m_i (\mathbf{v}_i - \mathbf{v}_m(y))^2, \quad (4)$$

quantifies the fluctuations around the mean velocity $\mathbf{v}_m(y) = 1/N(y) \sum_i^{N(y)} \mathbf{v}_i$.

The wind channel is divided along the y -direction in horizontal slices of volume $(2.5 \times 75 \times 7) D_{mean}^3$ to calculate the profiles of particle concentration, average particle velocity, alignment, and granular temperature. The particle concentration is the ratio between the volume of particles and the total volume. The flux profile is obtained for each slice from the product of the concentration and the average particle velocity.

Figure 3b shows the dependence of the saturated dimensionless flux (Eq. (3)) on e for different Shields numbers. For $\theta > 0.44$, the saturated flux is higher for $e = 0.65$ than for $e = 1.0$. The maximum at $e = 0.65$ increases substantially with the Shields number, i.e., the stronger the wind, the higher the peak. This is also confirmed by the flux profile in Fig. 4b, where the area below the curve, which corresponds to the total flux, is larger under dissipation. Interestingly, this optimal restitution coefficient is comparable to the values experimentally measured for collisions between grains of quartz sand [21]. Below we will explain this maximum in the flow at a specific value of the restitution coefficient as the result of the competition between two different mechanisms: the decreasing alignment of trajectories and the




FIG. 5. (color online) (a) The angle variance between particle trajectories for $e = 0.7$, $e = 1.0$, and without mid-air collisions, for $\theta = 0.9$. The inset shows the alignment a for the three cases. They confirm the weaker alignment in the simulations with elastic collisions. (b) Also for $\theta = 0.9$, the probability distribution for the maximum height, which can be fitted by the superposition of two exponential decays (solid (red) line): one for the leapers (dash-dotted (green) line) $[0.3 \exp(-0.12x)]$ and another for the saltons (dash-double-dotted (purple) line) $[0.0094 \exp(-0.0182x)]$, truncated for $y > 220$ possibly due to finite-size effects. For the sake of comparison, we include results without mid-air collisions (blue triangles), fitted by a single exponential decay (dashed (yellow) line) $[0.072 \exp(-0.0517x)]$. In the inset, we show the relation between the flying time and the maximum height, where the horizontal dashed line delimits the soft bed.

increasing uplift of particles with e .

In the absence of mid-air collisions, all particles are reptons (either creepers or leapers) and they follow stretched parabolic trajectories as typically assumed in saltation models [1]. With mid-air collisions, saltions emerge and they are sustained at higher positions, where the wind is stronger, by multiple collisions with leapers (as shown in Fig. 2). The number of collisions, and consequently the flying time of a salton, strongly depends on the concentration of leapers. Figure 3a compares the concentration profiles for $\theta = 0.90$ with and without mid-air collisions. The red (dark) curve obtained without collisions decays similarly as the one proposed by Jenkins and co-workers [22, 23]. In the presence of mid-air collisions (yellow (light) curve), particles are uplifted and, above a certain height, the concentration of particles is much larger than in the case without collisions.

Simultaneously, another important mechanism is activated. While, without dissipation, particles rebound randomly in the air, dissipation tends colliding particles to align their trajectories [24]. In the Aeolian transport, this symmetry breaking is expected to occur in the direction of the wind, affecting the overall flux. To investigate this alignment, we measure, for the dissipative ($e = 0.7$)




FIG. 6. (color online) Contribution to the total saturated flux of sheet flow (lower curve), the leapers in the soft bed (middle curve), and the saltions above the soft bed (upper curve) for $e = 0.7$ as function of the Shields number. The green area (lower region) shows the sheet flow in the particle bed $y < h_0$.

and the conservative ($e = 1.0$) cases, the granular temperature $T(y)$, the inverse of the temperature anisotropy given by $a(y) = \langle v_x^2(y) \rangle / T(y)$, and the variance of the velocity angle with respect to the wind direction, i.e., $\sigma_\alpha^2 = \langle \alpha^2 \rangle - \langle \alpha \rangle^2$, with $\alpha = \arctan(v_y/v_x)$. The granular temperature profile in Fig. 4a displays for $e = 1.0$ at every height a larger temperature for $e = 1.0$ as compared to $e = 0.7$ with a peak at a certain height. As a reference, at zero temperature, all particles would flow aligned with the same velocity. Figure 4c shows the dependence of the maximum temperature on the Shields number θ for both cases. The maximum temperature grows with $\theta^{1/2}$ (linear in u_*) for $e = 0.7$, and increases linearly with θ (quadratically with u_*) for $e = 1.0$, being always larger in the conservative case. This higher temperature could either be due to a larger dispersion in the magnitude or in the direction of the particle velocities. To distinguish these two possibilities we plot, in Fig. 5a, the variance of the velocity angle and the temperature anisotropy (main plot and inset, respectively). These data quantitatively confirm that the lower temperature in the inelastic case is mainly due to the alignment of particle trajectories. Clearly, the alignment of the particle trajectories, which tends to enhance the overall flux, decreases with e .

In the absence of mid-air collisions, the distribution of maximum heights can be approximated by a Poisson process and, therefore, well described by an exponential decay, as shown in Fig. 5b (dashed (black) line). With mid-air collisions, however, the distribution can only be described by the superposition of two exponentials. The first exponential dependence (dash-dotted (green) line) corresponds to the leapers, while the second one corresponds to the saltions, which typically fly above the

leapers (dash-dotted line). We can now quantitatively define the upper bound of the soft bed as the height above which the saltions are in majority. From the trajectories of individual particles we can also relate the flying time to the maximum height (see inset in Fig. 5b); we observe that saltions stay much longer in the air than leapers.

Three major types of transport contribute to the total flux in saltation: the sheet flow, the transport of leapers (in the soft bed), and of saltions (above the soft bed). Once identified the limits of the soft bed, for each Shields number, we can compute the contribution of each mechanism to the total flux, as shown in Fig. 6. The sheet flow, computed from the mass transport in the particle bed ($y < h_0$), results from the wind shear stress and the creep of particles on the surface. The contribution of the leapers and saltions is obtained from the flux in and above the soft bed, respectively. As observed in the figure, the relative contribution of the saltions and leapers significantly increases with the Shields number in comparison to the sheet flow. Figure 6 also shows that the saltions contribute the most to the total flux.

In the saturated flux, we also reproduced the discontinuity at the onset of saltation reported for 2D in Ref. [16]. We also studied the impact of mid-air collisions in two dimensions. The decrease in the spatial dimension enhances the relevance of mid-air collisions and the consequences discussed here are even more pronounced. However, as expected, the curves for the saturated flux in 2D and 3D overlap if the Shields number in 2D is rescaled by an appropriate factor that takes into account the width of the system.

Summarizing, the contribution of mid-air collisions cannot be neglected in saltation. The absence of collisions strongly underestimate the total mass flux. The saltions contribute significantly to the flux enhancement by acquiring large momentum from the wind and using it to eject more particles through more intense splashes. We provide a new picture of Aeolian saltation where the competition between uplift due to mid-air collisions and alignment due to inelasticity optimizes the mass transport in the presence of dissipative collisions. Interestingly, the position of the maximum corresponds to the restitution coefficient typically observed in granular collisions.

Our results are crucial for future studies in Aeolian saltation, since they provide qualitative and quantitative information about the influence of mid-air collisions, which should be considered in future modeling. Future work might include the aerodynamic lift of the particles, turbulent wind speed fluctuations, and the electrostatic interaction between particles giving a contribution of long range forces.

This work has been supported by the National Council for Scientific and Technological Development CNPq, Brazil and ETH Grant (No. ETH-10 09-2). We acknowledge also the financial support from the

European Research Council (ERC) Advanced Grant 319968-FlowCCS, Natural Science Foundation of China 51009120, Zhejiang Province Ocean and Fisheries Bureau 2010210, and Zhejiang University 2012HY012B.

* cmarcus@ethz.ch

† nuno@ethz.ch

‡ tpaeltz@ethz.ch

§ hans@ifb.baug.ethz.ch

- [1] M. P. Almeida, J. S. Andrade, and H. J. Herrmann, Phys. Rev. Lett. **96**, 21 (2006).
- [2] M. P. Almeida, E. J. R. Parteli, J. S. Andrade, and H. J. Herrmann, Proc. Natl. Acad. Sci. USA **105**, 6222 (2008).
- [3] B. T. Werner and P. K. Haff, Sedimentology **35**, 189 (1988).
- [4] I. K. McEwan and B. B. Willetts, J. Fluid Mech. **252**, 99 (1993).
- [5] J. F. Kok and N. O. Renno, J. Geophys. Res. **114**, 149 (2009).
- [6] T. Pähtz, J. Kok, and H. J. Herrmann, New J. Phys. **14**, 43035 (2012).
- [7] R. A. Bagnold, *The Physics of Blown Sand and Desert Dunes* (Methuen, New York, 1941) p. 265.
- [8] J. F. Kok, E. R. Parteli, T. Michaels, and D. B. Karam, Rep. Prog. Phys. **75**, 106901 (2012).
- [9] S. H. Williams, *A comparative planetological study of particle speed and concentration during aeolian saltation*, Ph.D. thesis, Arizona State University (1987), ph.D thesis.
- [10] M. Soerensen and I. McEwan, Sedimentology **43**, 65 (1996).
- [11] B. Andreotti, J. Fluid Mech. **510**, 47 (2004).
- [12] M. Lämmel, D. Rings, and K. Kroy, New J. Phys. **14**, 093037 (2012).
- [13] S. Mitha, M. Q. Tran, B. T. Werner, and P. K. Haff, Acta Mechanica **63**, 267 (1986).
- [14] R. S. Anderson and P. K. Haff, Science **241**, 820 (1988).
- [15] M. Griebel, S. Knapek, and G. Zumbusch, *Numerical Simulation in Molecular Dynamics: Numerics, Algorithms, Parallelization, Applications* (Springer, New York, 2007).
- [16] M. V. Carneiro, T. Pähtz, and H. J. Herrmann, Phys. Rev. Lett. **107**, 098001 (2011).
- [17] See Supplemental Material at.
- [18] F. Rioual, A. Valance, and D. Bideau, Phys. Rev. E **62**, 2450 (2000).
- [19] F. Rioual, Ph.D. thesis, University of Rennes 1 (2002).
- [20] F. Rioual, A. Valance, and D. Bideau, Europhys. Lett. **61**, 194 (2003).
- [21] M. Banks, N. Bridges, and M. Benzit, Proceedings of the 36th Annual Lunar and Planetary Science Conference , 2116 (2005).
- [22] J. T. Jenkins, I. Cantat, and A. Valance, Phys. Rev. E **82**, 020301R (2010).
- [23] J. M. Pasini and J. T. Jenkins, Phil. Trans. R. Soc. A **363**, 1625 (2005).
- [24] Y. L. Duparcmeur, H. Herrmann, and J. Troadec, J. Physique I **5**, 1119 (1995).

Supplemental Material: Mid-air collisions enhance saltation

M. V. Carneiro¹, N. A. M. Araújo¹, T. Pähtz^{2,3}, and H. J. Herrmann^{1,4}

¹*Institut für Baustoffe, ETH-Hönggerberg, Schafmattstrasse 6, 8093 Zürich, Switzerland*

²*Department of Ocean Science and Engineering, Zhejiang University, 310058 Hangzhou, China*

³*State Key Laboratory of Satellite Ocean Environment Dynamics,
Second Institute of Oceanography, Hangzhou, China and*

⁴*Departamento de Física, Universidade Federal do Ceará, 60451-970 Fortaleza, Ceará, Brazil*

(Dated: December 18, 2012)

I. PARTICLE DYNAMICS

Trajectories and velocities of the particles are obtained iteratively by solving the Newton equations of motion, through the velocity-Störmer-Verlet scheme [1]. The contact between particles is described by the spring dashpot potential, which has an elastic and a dissipative contribution acting on each particle. When two particles i and j overlap (i.e. when their distance is smaller than the sum of their radii) an elastic force is applied,

$$\mathbf{F}_{el}^{(i)} = km_i[|\mathbf{r}_{ij}| - 1/2(d_i + d_j)] \frac{\mathbf{r}_{ij}}{|\mathbf{r}_{ij}|}, \quad (1)$$

where $k = 0.5$ is a spring constant, d_i and d_j are the diameters, m_i is the mass of particle i , and $\mathbf{r}_{ij}$ points from particle i to j . A dissipation force is also applied accounting for the inelasticity of the collision,

$$\mathbf{F}_{diss}^{(i)} = -\gamma \mathbf{v}_{ij}, \quad (2)$$

where $\mathbf{v}_{ij} = \mathbf{v}_i - \mathbf{v}_j$ is the relative velocity and γ is the dissipation coefficient. The coefficient of restitution e is given by the ratio between the absolute velocities after and before the collision. For particle-wall collisions, the same forces act as if particle i would collide with another particle of diameter d_i [2]. For simplicity, friction and rotation are neglected.

II. THE WIND PROFILE

A logarithmic velocity field mimics the wind profile in the absence of saltating grains in the x -direction with

$$u(y) = \frac{u_*}{\kappa} \ln \frac{y - h_0}{y_0}, \quad (3)$$

where $y_0 = D_{mean}/30$ is the roughness of the bed with D_{mean} the mean diameter of the particles, h_0 the bed height, $\kappa = 0.4$ the von Kármán constant, and u_* the wind shear velocity. In the presence of grains, the wind strength is substantially reduced due to the momentum transferred to the grains [3]. The grain stress $\tau_g(y)$ quantifies the average horizontal force per unit area f that the wind applies on the grains above y [4], i.e.,

$$\tau_g(y) = \int_y^\infty f(y') dy'. \quad (4)$$

where f is the average horizontal force per unit volume. The modified wind shear velocity $u_\tau(y)$ is the fluid stress left at the height y after the momentum transfer,

$$u_\tau(y) = u_* \sqrt{1 - \frac{\tau_g}{\rho_w u_*^2}}, \quad (5)$$

where ρ_w is the air density. To obtain the modified wind profile [5], we solve,

$$\frac{du}{dy} = \frac{u_\tau(y)}{\kappa y}, \quad (6)$$

considering $u_\tau(y)$ constant within an interval dy .

The numerical solution is achieved iteratively. The position of the bed surface is used as starting point of the integration of the wind profile. However, the particle splash and the sheet flow dynamically affect the shape of the bed surface. Consequently, we need to find h_0 for every time step. We use a criterion based on the wind values to define an approximate position for the particle surface. High density areas strongly reduce wind velocities. If the calculated velocity v_i in area y_i is below $0.1u_*$, we assume that this area contains the particle bed and the velocity is set to zero. This means that h_0 is chosen to be the point where the calculated velocity exceeds $0.1u_*$.

The wind drag is the only external force applied to the particles along the x -direction,

$$\mathbf{F}_d = -\frac{\pi D^2}{8} \rho_w C_d v_r \mathbf{v}_r, \quad (7)$$

where ρ_w is the air density and $\mathbf{v}_r = \mathbf{v} - \mathbf{u}$ is the velocity difference between particle and wind, with $v_r = |\mathbf{v}_r|$. The drag coefficient C_d proposed by Cheng [6] is suited to model grains with irregular, natural, shapes, and is given by,

$$C_d = \left[\left(\frac{32}{Re} \right)^{2/3} + 1 \right]^{3/2}, \quad Re = \frac{\rho_w v_r D_{mean}}{\mu}, \quad (8)$$

where $\mu = 1.8702 \times 10^{-5}$ kg/(m.s) is the dynamic viscosity and Re , the Reynolds number.

-
- [1] M. Griebel, S. Knapek, and G. Zumbusch, *Numerical Simulation in Molecular Dynamics: Numerics, Algorithms, Parallelization, Applications* (Springer, 2007).
 - [2] H. Herrmann, *Physica A* **191**, 236 (1992).
 - [3] R. S. Anderson and P. K. Haff, *Science* **241(4867)**, 820 (1988).
 - [4] J. E. Ungar and P. Haff, *Sedimentology* **34(2)**, 289 (1987).
 - [5] R. S. Anderson and P. K. Haff, *Acta Mech.* **1**, 21 (1991).
 - [6] N. S. Cheng, *J. Hydraul. Eng.* **123**, 149 (1997).

